

Prednisolone

Prednisolone is a steroid medication.

Its side effects include

1. Heartburn, gastric discomfort or formation of ulcers in the stomach.
2. Hip pain (rarely death of the top part of the hip bone).
3. Fluid retention
4. Mood alteration.
5. Destabilisation of diabetes control.

“Do’s”

Always take medication with food.

Notify the doctor who has prescribed the medication or your local doctor if you experience any of the above side effects.

“Do not’s”


Do not take Prednisolone if you are already taking a blood thinning medication such as Warfarin or Coumadin.

Do not continue to use the medication if you experience any of the above side effects.

Do not use Aspirin, anti-inflammatory drugs i.e. Voltaren, Brufen, Indocid, etc while taking Prednisolone.

- Paracetamol and Codeine **are safe** to take for pain while on Prednisolone.

Do not drink alcohol while taking Prednisolone.



North Melbourne ENT

ABN 28 582 450 307

100/30 Wreckyn Street
North Melbourne VIC 3051
T 03 9078 8074
F 03 9078 8105
northmelbourne@nment.com.au
www.nment.com.au

Also located at:

Williamstown ENT

54 Electra Street
Williamstown VIC 3016
T 03 9397 5507
F 03 9397 6914
williamstown@nment.com.au

A/Prof Tim Iseli

MBBS (hons.), FRACS

ENT Surgeon

Provider No. 231248MT (N)

Provider No. 231248NB (W)

Dr Claire Iseli

MBBS (hons.), FRACS, MS

ENT Surgeon

Provider No. 246559JX (N)

Provider No. 246559PB (W)

*An association of
independent practitioners*